

Linear theory of weakly amplified, parallel propagating, transverse temperature anisotropy instabilities in magnetized thermal plasmas

R. Schlickeiser^{1,2}. T. Skoda¹

1 Institut für Theoretische Physik, Lehrstuhl IV: Weltraum- und Astrophysik,
Ruhr-Universität Bochum, D-44780 Bochum, Germany

2 Research Department Plasmas with Complex Interactions, Ruhr-Universität Bochum,
D-44780 Bochum, Germany

rsch@tp4.rub.de, ts@tp4.rub.de

Received 2010 February 1; accepted _____

ABSTRACT

A rigorous analytical study of the dispersion relations of weakly amplified transverse fluctuations with wave vectors ($\vec{k} \parallel \vec{B}$) parallel to the uniform background magnetic field $\vec{B}$ in an anisotropic bi-Maxwellian magnetized electron-proton plasma is presented. A general analytical instability condition is derived that holds for different values of the electron (A_e) and proton (A_p) temperature anisotropies. We determine the conditions for which the weakly amplified LH-handed polarized Alfvén-proton-cyclotron and RH-handed polarized Alfvén-Whistler-electron-cyclotron branches can be excited. For different regimes of the electron plasma frequency phase speed $w = \omega_{p,e}/(kc)$ these branches reduce to the RH and LH polarized Alfvén waves, RH polarized high- and low-phase speed Whistler, RH polarized proton and LH polarized electron cyclotron modes. Analytic instability threshold conditions are derived in terms of the combined temperature anisotropy $A = T_{\perp}/T_{\parallel}$, the parallel plasma beta $\beta_{\parallel} = 8\pi n_e k_B T_{\parallel}/B^2$ and the electron plasma frequency phase speed $w = \omega_{p,e}/(kc)$ for each mode.

The results of our instability study are applied to the observed solar wind magnetic turbulence at values of $90 \leq w \leq 330$. According to the existence conditions of the different instabilities, only the left-handed and right-handed polarized Alfvén wave instabilities can operate here. Besides the electron-proton mass ratio $\mu = 1836$, the Alfvénic instability threshold conditions are controlled by the single observed plasma parameter w . The Alfvénic instability diagram explains well the main characteristic properties of the observed solar wind fluctuations. Especially, the observed confinement limits at small parallel plasma beta values are explained.

Subject headings: plasmas – instabilities – turbulence – magnetic fields – solar wind

1. Introduction

Kinetic plasma relaxation and turbulence generation processes are responsible for the observed properties of the solar wind plasma which is the only cosmic collisionless plasma accessible to detailed in-situ satellite observations (Bale et al. 2009). Although the detailed plasma relaxation processes are not understood, the observed electron and proton distribution functions are close to bi-Maxwellian distributions with different temperatures along and perpendicular to the ordered magnetic field direction. Ten years of Wind/SWE data (Kasper et al. 2002) have demonstrated that the proton and electron temperature anisotropies $A = T_{\perp}/T_{\parallel}$ are bounded by mirror and firehose instabilities (Hellinger et al. 2006) at large values of the parallel plasma beta $\beta_{\parallel} = 8\pi n k_B T_{\parallel}/B^2 \geq 1$. In the parameter plane defined by the temperature anisotropy $A = T_{\perp}/T_{\parallel}$ and the parallel plasma beta $\beta_{\parallel}$, stable plasma configuration are only possible within a rhomb-like configuration around $\beta_{\parallel} \simeq 1$, whose limits are defined by the threshold conditions for the mirror and firehose instabilities. If a plasma would start with parameter values outside this rhomb-like configuration, it immediately would generate fluctuations via the mirror and firehose instabilities, which quickly relax the plasma distribution into the stable regime within the rhomb-configuration. The plasma parameters of other dilute cosmic plasmas including the interstellar and intracluster medium (Schekochihin et al. 2005) and accretion disks around compact, massive objects (Sharma et al. 2007) are similar to the solar wind plasma, so that the temperature-anisotropy instabilities should also operate in these systems.

The study of linear electromagnetic instabilities in a collisionless, homogeneous, magnetized, electron-proton plasma with temperature anisotropies has a long history; for reviews we refer the interested reader to Ch. 7 of the monograph by Gary (1993), Cupperman (1992) and Marsch (2006). Hellinger et al. (2006) considered linear instability calculations for the oblique mirror, firehose and proton cyclotron instabilities, and represented the

approximate threshold conditions from the work of Gary et al. (1994, 2001), Gary & Lee (1994), Samsonov et al. (2001) and Pokhotelov et al. (2004) by analytic relations of the form $A = 1 + a(\beta_{\parallel} - \beta_0)^{-b}$ where a , b and β_0 are fitted parameter values.

In order to understand the confinement limits also at small values of the parallel plasma beta $\beta_{\parallel} < 1$, we analyze here rigorously the full linear dispersion relation in a collisionless homogenous plasma with anisotropic ($A \neq 1$) bi-Maxwellian particle velocity distributions of electrons and protons for electromagnetic fluctuations with wave vectors ($\vec{k} \times \vec{B} = 0$) parallel to the uniform background magnetic field $\vec{B}$. With a typical solar wind temperature $T = 10^5 T_5 \text{K}$, the thermal particle energy $k_B T = 8.6 T_5 \text{ eV}$ is much less than the electron rest mass $m_e c^2 = 5.11 \cdot 10^5 \text{ eV}$, so that the use of the nonrelativistic linearized Vlasov/Maxwell equations is justified. For weakly ($\gamma \ll \omega_R$) amplified wave solutions we derive analytically existence and instability conditions. ω_R and γ here refer to real and imaginary part of the complex frequency $\omega = \omega_R + i\gamma$. Our analysis follows closely the recent study (Schlickeiser 2010) for equal-mass pair plasmas – hereafter referred to as paper S. Our investigation is restricted to weakly amplified ($\gamma \ll \omega_R$) solutions covering the left-handed (LH) polarized Alfvén-proton cyclotron branch and the right-handed (RH) polarized Alfvén-Whistler-electron cyclotron branch. The corresponding analysis of weakly propagating ($\omega_R \ll \gamma$) solutions, including mirror, firehose, electron cyclotron and cold magnetized Weibel fluctuations, is the subject of a subsequent paper. As we will demonstrate below, for the case of parallel ($\vec{k} \times \vec{B} = 0$) wavevectors simple analytical threshold conditions for the two branches can be derived in terms of the combined temperature anisotropy $A = T_{\perp}/T_{\parallel}$, the parallel plasma beta $\beta_{\parallel} = 8\pi n_e k_B T_{\parallel}/B^2$, the electron-proton mass ratio and the electron plasma frequency phase speed $w = \omega_{p,e}/(kc)$.

2. Dispersion relations

2.1. Basic equations

For a nonzero background magnetic field strength the nonrelativistic dispersion relations for right-handed (RH) and left-handed (LH) polarized fluctuations with wave vectors $\vec{k} \times \vec{B} = 0$ in a thermal electron-proton plasma are (Gary 1993)

$$0 = D_{RH,LH}(k, \omega) = \omega^2 - k^2 c^2 + \sum_{a=p,e} \omega_{p,a}^2 \left[\frac{\omega}{\sqrt{2} k u_{a,\parallel}} Z \left(\frac{\omega \pm \Omega_a}{\sqrt{2} k u_{a,\parallel}} \right) + \frac{1}{2} (1 - A_a) Z' \left(\frac{\omega \pm \Omega_a}{\sqrt{2} k u_{a,\parallel}} \right) \right] = 0, \quad (1)$$

where we sum over a proton (p)-electron (e) plasma. and where $k = |k_{\parallel}|$. $\omega_{p,e}$ denotes the electron plasma frequency, $u_{a,\parallel} = (k_B T_{a,\parallel}/m_a)^{1/2}$ is the parallel thermal velocity of component a , $\Omega_a = e_a B/(m_a c)$ is the non-relativistic gyrofrequency, and $A_a = T_{a,\perp}/T_{a,\parallel}$ is the temperature anisotropy of component a , where the directional subscripts refer to directions relative to the background magnetic field. The dispersion relations (1) allow for different values of the proton and electron parallel temperatures and temperature anisotropies.

$Z(x)$ and $Z'(x)$ denote the plasma dispersion function (Fried & Conte 1961) and its derivative

$$Z(x) = \pi^{-1/2} \int_{-\infty}^{\infty} dt \frac{e^{-t^2}}{t - x} \quad (2)$$

with the well-known properties

$$Z'(x) = -2 [1 + xZ(x)], \quad (3)$$

and

$$Z(-x) = 2\pi^{1/2}\imath e^{-x^2} - Z(x), \quad Z'(-x) = 4\pi^{1/2}\imath x e^{-x^2} + Z'(x) \quad (4)$$

We will frequently use the asymptotic expansions

$$Z(x) \simeq \imath\pi^{1/2}e^{-x^2} - 2x \left[1 - \frac{2x^2}{3} \right], \quad |x| \ll 1 \quad (5)$$

and

$$Z(x) \simeq \imath\sigma\pi^{1/2}e^{-x^2} - \frac{1}{x} \left[1 + \frac{1}{2x^2} + \frac{3}{4x^4} \right], \quad |x| \gg 1 \quad (6)$$

where $\sigma = 0$ if $\Im(x) > 0$, $\sigma = 1$ if $\Im(x) = 0$ and $\sigma = 2$ if $\Im(x) < 0$.

Paper S has demonstrated that the analytical analysis is enormously faoilitated if we work with space speed rather than frequencies. We therefore introduce the complex phase speeds

$$f = \frac{\omega}{kc} = \frac{\omega_R + \imath\gamma}{kc} = R + \imath S, \quad R = \frac{\omega_R}{kc}, \quad S = \frac{\gamma}{kc}, \quad (7)$$

the plasma frequency phase speed

$$w = \frac{\omega_{p,e}}{kc}, \quad (8)$$

and the absolute value of the electron gyrofrequency phase speed

$$b = \frac{|\Omega_e|}{kc}, \quad (9)$$

where $|\Omega_e| = eB/m_e c$ is the absolute value of the electron gyrofrequency. We also introduce the mass ratio

$$\mu = m_p/m_e = 1836, \quad (10)$$

and the dimensionless proton and electron temperatures

$$\Theta_e \equiv \left(\frac{2k_B T_{e,\parallel}}{m_e c^2} \right)^{1/2}, \quad \Theta_p \equiv \left(\frac{2k_B T_{p,\parallel}}{m_p c^2} \right)^{1/2} \quad (11)$$

Throughout this work, in the classification of Swanson (1989) we discuss high density plasmas with $\omega_{p,e} \gg |\Omega_e|$, corresponding to $w \gg b$ which applies to nearly all astrophysical plasmas. These plasmas are dense enough that the electron plasma frequency is much larger than the electron gyrofrequency, but small enough that elastic Coulomb collisions can be neglected.

The two dispersion relations (1) then read

$$\begin{aligned} 0 = \frac{D_{RH,LH}(k, f)}{k^2 c^2} = \Lambda_{RH,LH}(k, f) = f^2 - 1 + \\ \frac{w^2}{\mu} \left[\frac{f}{\Theta_p} Z \left(\frac{f \pm \frac{b}{\mu}}{\Theta_p} \right) + \frac{1}{2} (1 - A_p) Z' \left(\frac{f \pm \frac{b}{\mu}}{\Theta_p} \right) \right] \\ + w^2 \left[\frac{f}{\Theta_e} Z \left(\frac{f \mp b}{\Theta_e} \right) + \frac{1}{2} (1 - A_e) Z' \left(\frac{f \mp b}{\Theta_e} \right) \right] \end{aligned} \quad (12)$$

We notice the symmetry $\Lambda(-k_{\parallel}, f) = \Lambda(k_{\parallel}, f)$ of both dispersion relations allowing to consider only positive values of the wavenumber $k > 0$. In the following we will simplify the analysis by considering only equal parallel temperature plasmas ($T_{e,\parallel} = T_{p,\parallel}$) so that $\Theta_e = \Theta$ and $\Theta_p = \Theta/\mu^{1/2}$.

The dispersion relations (12) can be separated into real and imaginary parts $\Lambda(R, S) = \Re \Lambda(R, S) + i \Im \Lambda(R, S) = 0$, implying the two conditions

$$\Re \Lambda(R, S) = 0, \quad \Im \Lambda(R, S) = 0 \quad (13)$$

In terms of the complex phase speed $f = R + \imath S$ the real and imaginary parts of the two dispersion relations (12) read

$$\begin{aligned}
0 = \Re \Lambda_{RH,LH}(R, S) = R^2 - S^2 - 1 + \\
w^2 \left[\frac{R}{\Theta \mu^{1/2}} \Re Z \left(\frac{\mu^{1/2}}{\Theta} \left[R + \imath S \pm \frac{b}{\mu} \right] \right) + \frac{1 - A_p}{2\mu} \Re Z' \left(\frac{\mu^{1/2}}{\Theta} \left[R + \imath S \pm \frac{b}{\mu} \right] \right) \right. \\
\left. + \frac{R}{\Theta} \Re Z \left(\frac{R + \imath S \mp b}{\Theta} \right) + \frac{1 - A_e}{2} \Re Z' \left(\frac{R + \imath S \mp b}{\Theta} \right) \right] \\
- \frac{w^2 S}{\Theta} \left[\frac{1}{\mu^{1/2}} \Im Z \left(\frac{\mu^{1/2}}{\Theta} \left[R + \imath S \pm \frac{b}{\mu} \right] \right) + \Im Z \left(\frac{R + \imath S \mp b}{\Theta} \right) \right]
\end{aligned} \tag{14}$$

and

$$\begin{aligned}
0 = \Im \Lambda_{RH,LH}(R, S) = 2RS + \frac{w^2}{2} \left[\frac{1 - A_p}{\mu} \Im Z' \left(\frac{\mu^{1/2}}{\Theta} \left[R + \imath S \pm \frac{b}{\mu} \right] \right) \right. \\
\left. + (1 - A_e) \Im Z' \left(\frac{R + \imath S \mp b}{\Theta} \right) \right] \\
+ \frac{w^2 S}{\Theta} \left[\frac{1}{\mu^{1/2}} \Re Z \left(\frac{\mu^{1/2}}{\Theta} \left[R + \imath S \pm \frac{b}{\mu} \right] \right) + \Re Z \left(\frac{R + \imath S \mp b}{\Theta} \right) \right] \\
+ \frac{w^2 R}{\Theta} \left[\frac{1}{\mu^{1/2}} \Im Z \left(\frac{\mu^{1/2}}{\Theta} \left[R + \imath S \pm \frac{b}{\mu} \right] \right) + \Im Z \left(\frac{R + \imath S \mp b}{\Theta} \right) \right]
\end{aligned} \tag{15}$$

Here we consider solutions of the dispersion relations in the weak damping/amplification limit $|S| \ll R$. As shown in paper S in this limit the real part of the dispersion relation satisfies

$$\Re \Lambda(R, S = 0) = 0, \tag{16}$$

whereas the corresponding imaginary part is given by

$$S = - \frac{\Im \Lambda(R, S = 0)}{\frac{\partial \Re \Lambda(R, S=0)}{\partial R}} \tag{17}$$

As before we introduce the parallel plasma beta

$$\beta_{\parallel} = \frac{\Theta^2 w^2}{b^2} = \frac{8\pi n_e k_B T_{\parallel}}{B^2}, \quad (18)$$

which expresses the magnetic field strength as

$$b = \frac{\Theta w}{\beta_{\parallel}^{1/2}} \quad (19)$$

in terms of the electron plasma frequency (w), the parallel temperature (Θ) and the parallel plasma beta $\beta_{\parallel}$. Our restriction to high-density plasmas with $\omega_{p,e} > |\Omega_e|$ or $w > b$ then requires to have parallel plasma betas $\beta_{\parallel} > \Theta^2$.

3. Weakly damped and amplified solutions

For weakly damped or amplified fluctuations Eqs. (14) – (15) read

$$\begin{aligned} 0 = \Re \Lambda_{RH,LH}(R, S = 0) = R^2 - 1 \\ + w^2 \left[\frac{R}{\Theta} \Re Z \left(\frac{R \mp b}{\Theta} \right) + \frac{1 - A_e}{2} \Re Z' \left(\frac{R \mp b}{\Theta} \right) \right. \\ \left. + \frac{R}{\Theta \mu^{1/2}} \Re Z \left(\frac{\mu^{1/2}}{\Theta} \left[R \pm \frac{b}{\mu} \right] \right) + \frac{1 - A_p}{2\mu} \Re Z' \left(\frac{\mu^{1/2}}{\Theta} \left[R \pm \frac{b}{\mu} \right] \right) \right] \end{aligned} \quad (20)$$

and

$$\begin{aligned} \Im \Lambda_{RH,LH}(R, S = 0) = \frac{w^2}{2} \left[\frac{1 - A_p}{\mu} \Im Z' \left(\frac{\mu^{1/2}}{\Theta} \left[R \pm \frac{b}{\mu} \right] \right) + (1 - A_e) \Im Z' \left(\frac{R \mp b}{\Theta} \right) \right] \\ + \frac{w^2 R}{\Theta} \left[\frac{1}{\mu^{1/2}} \Im Z \left(\frac{\mu^{1/2}}{\Theta} \left[R \pm \frac{b}{\mu} \right] \right) + \Im Z \left(\frac{R \mp b}{\Theta} \right) \right], \end{aligned} \quad (21)$$

which have to be investigated for positive values of $R \geq 0$.

As noted before, these dispersion relations can be further reduced with the asymptotic expansions (5) and (6), depending on the absolute values of the arguments

$$P_{\pm}(R) = \frac{\mu^{1/2}}{\Theta} \left[\left| R \pm \frac{b}{\mu} \right| \right], \quad E_{\pm}(R) = \frac{|R \pm b|}{\Theta} \quad (22)$$

of the plasma dispersion function and its derivative being small or large compared to unity. Scaling $R = bx$, corresponding to $x = \omega_R/|\Omega_e|$, the absolute values of the arguments (22) in terms of the parallel plasma beta read

$$P_{\pm}(x) = \frac{w\mu^{1/2}}{\beta_{\parallel}^{1/2}} \left[\left| x \pm \frac{1}{\mu} \right| \right], \quad E_{\pm}(x) = \frac{w}{\beta_{\parallel}^{1/2}} |x \pm 1|, \quad (23)$$

which are shown in Fig. 1 as a function of the normalized real frequency x .

For parallel plasma beta values $\beta_{\parallel} \ll (w^2/\mu)$ we note that $P_+(x) \gg 1$, $E_+(x) \gg 1$ for all values of x , whereas $P_-(x) \gg 1$ for x outside the small interval

$$x \notin \left[\frac{1}{\mu} - \frac{\beta_{\parallel}^{1/2}}{w\mu^{1/2}}, \frac{1}{\mu} + \frac{\beta_{\parallel}^{1/2}}{w\mu^{1/2}} \right] \quad (24)$$

around the proton cyclotron frequency, and $E_-(x) \gg 1$ for x outside the small interval

$$x \notin \left[1 - \frac{\beta_{\parallel}^{1/2}}{w}, 1 + \frac{\beta_{\parallel}^{1/2}}{w} \right] \quad (25)$$

around the electron cyclotron frequency.

In the following we limit our analysis to such values of the parallel plasma beta $\beta_{\parallel} \ll (w^2/\mu)$, so that the asymptotic expansion (6) of the plasma dispersion function can be used except near the indicated proton and electron cyclotron frequencies.

Irrespective of the values of $P_{\pm}$ and $E_{\pm}$ we note that both asymptotic expansions (5) and (6) yield the same imaginary part of the dispersion relation




Fig. 1.— Absolute values of the arguments $P_{\pm}(x)$ and $E_{\pm}(x)$ of the plasma dispersion function and its derivative in the weak damping/amplification limit as a function of $x = R/b$.

$$\begin{aligned}
 0 = \Im \Lambda_{RH,LH}(R, S = 0) &= \pi^{1/2} w^2 \frac{b}{\Theta} \left(\frac{1}{\mu^{1/2}} \left[A_p \left[\frac{R}{b} \pm \frac{1}{\mu} \right] \mp \frac{1}{\mu} \right] e^{-\frac{\mu}{\Theta^2} (R \pm \frac{b}{\mu})^2} \right. \\
 &\quad \left. + \left[A_e \left[\frac{R}{b} \mp 1 \right] \pm 1 \right] e^{-\frac{(R-b)^2}{\Theta^2}} \right), \tag{26}
 \end{aligned}$$

4. Alfven, Whistler, cyclotron waves and electromagnetic light

For parallel plasma beta values $\beta_{\parallel} \ll (w^2/\mu)$ the asymptotic expansion (6) yields

$$\Re Z \left(\frac{R \pm b}{\Theta} \right) \simeq -\frac{\Theta}{R \pm b} \left[1 + \frac{\Theta^2}{2(R \pm b)^2} \right], \quad (27)$$

$$\Re Z' \left(\frac{R \pm b}{\Theta} \right) \simeq +\frac{\Theta^2}{(R \pm b)^2} \left[1 + \frac{3\Theta^2}{2(R \pm b)^2} \right], \quad (28)$$

$$\Re Z \left(\frac{\mu^{1/2}}{\Theta} \left[R \pm \frac{b}{\mu} \right] \right) \simeq -\frac{\Theta}{\mu^{1/2}(R \pm \frac{b}{\mu})} \left[1 + \frac{\Theta^2}{2\mu(R \pm \frac{b}{\mu})^2} \right] \quad (29)$$

and

$$\Re Z' \left(\frac{\mu^{1/2}}{\Theta} \left[R \pm \frac{b}{\mu} \right] \right) \simeq +\frac{\Theta^2}{\mu(R \pm \frac{b}{\mu})^2} \left[1 + \frac{3\Theta^2}{2\mu(R \pm \frac{b}{\mu})^2} \right] \quad (30)$$

We then obtain for the real part of the dispersion relations (20) to lowest order in $\Theta^2 \ll 1$

$$\begin{aligned} 0 = \Re \Lambda_{RH,LH}(R, S = 0) &= R^2 - 1 - w^2 R \left[\frac{1}{R \mp b} + \frac{1}{\mu(R \pm \frac{b}{\mu})} \right] \\ &+ \frac{\Theta^2 w^2}{2} \left[\frac{1 - A_e}{(R \mp b)^2} - \frac{R}{(R \mp b)^3} \right] + \frac{\Theta^2 w^2}{2\mu^2} \left[\frac{1 - A_p}{(R \pm \frac{b}{\mu})^2} - \frac{R}{(R \pm \frac{b}{\mu})^3} \right] \end{aligned} \quad (31)$$

With the scaling $R = bx$ the dispersion relation (31) reads

$$\begin{aligned} 0 = \Re \Lambda_{RH,LH}(x, S = 0) &= \left[b^2 + \frac{(1 + \mu)w^2}{(1 \pm \mu x)(1 \mp x)} \right] x^2 - 1 \\ &\pm \frac{\beta_{\parallel}}{2} \left(\frac{A_e x \pm (1 - A_e)}{(1 \mp x)^3} - \frac{A_p \mu x \mp (1 - A_p)}{(1 \pm \mu x)^3} \right) \end{aligned} \quad (32)$$

where we introduce the parallel plasma beta (18).

In different limits the solutions of Eqs. (31) and (32) describe Alfvén waves, Whistler waves, cyclotron waves and electromagnetic light. We consider each case in the next sections.

4.1. Alfven waves at phase speeds $R \ll b/\mu$ for $\Theta^2 \ll \beta_{\parallel} < w^2/\mu$

For phase speeds $R \ll b/\mu$ the dispersion relation (31) simplifies to

$$\begin{aligned} 0 = \Re \Lambda_{RH,LH}(R, S = 0) &= R^2 \left(1 + \frac{2w^2(1+\mu)}{b^2} \right) - 1 \\ &\quad + \frac{\beta_{\parallel}}{2} \left[2 - A_e - A_p \pm \frac{2R}{b}(3 - A_e) \mp \frac{2\mu R}{b}(3 - A_p) \right] \\ &\simeq R^2 \left(1 + \frac{c^2}{V_A^2} \right) - [1 + (A - 1)\beta_{\parallel}], \end{aligned} \quad (33)$$

where we introduce the Alfven speed

$$\frac{(1+\mu)w^2}{b^2} = \frac{4\pi n_e(m_e + m_p)c^2}{B^2} = \frac{c^2}{V_A^2}, \quad (34)$$

the parallel plasma beta (18) and the combined plasma temperature anisotropy

$$A = \frac{A_p + A_e}{2} \quad (35)$$

For high-density plasmas $V_A \ll c$, the dispersion relation (33) yields the LH and RH polarized Alfven modes with the same

$$R \simeq \frac{b}{\sqrt{1+\mu}w} \sqrt{1 + (A-1)\beta_{\parallel}} = \Theta \sqrt{\frac{1 + (A-1)\beta_{\parallel}}{(1+\mu)\beta_{\parallel}}} = \frac{V_A}{c} \sqrt{1 + (A-1)\beta_{\parallel}}, \quad (36)$$

which either can propagate forward and backward. Note that the condition $R \ll b/\mu$ requires $(w^2/\mu) \gg 1$.

Moreover, the four Alfven modes only exist for temperature anisotropies such that $1 + (A-1)\beta_{\parallel} \geq 0$ corresponding to

$$A > \left(1 - \frac{1}{\beta_{\parallel}}\right) \quad (37)$$

which includes the isotropic ($A_p = A_e = A = 1$) plasma temperature case. For small plasma betas ($\beta_{\parallel} \leq 1$) the condition (37) is always fulfilled whereas for large plasma betas ($\beta_{\parallel} > 1$) the combined anisotropy A has to be larger than $1 - \beta_{\parallel}^{-1}$.

Eq. (33) also provides

$$\frac{\partial \Re \Lambda_{RH,LH}(R, S=0)}{\partial R} \simeq 2 \frac{c^2}{V_A^2} R, \quad (38)$$

so that for all four modes according to Eqs. (17) and (26) the growth/damping rate is

$$S_{RH,LH} = -\frac{V_A^2}{2c^2 R} \Im \Lambda_{RH,LH}(R, S=0) = -\frac{\pi^{1/2} b^3}{2(1+\mu)\Theta R} \left(\frac{1}{\mu^{1/2}} \left[A_p \left[\frac{R}{b} \pm \frac{1}{\mu} \right] \mp \frac{1}{\mu} \right] e^{-\frac{\mu}{\Theta^2} (R \pm \frac{b}{\mu})^2} + \left[A_e \left[\frac{R}{b} \mp 1 \right] \pm 1 \right] e^{-\frac{(R \mp b)^2}{\Theta}} \right) \quad (39)$$

For isotropic ($A_p = A_e = 1$) plasma temperatures all four Alfvén modes are damped in agreement with the general theorem of Brinca (1990) on the electromagnetic stability of isotropic plasma populations. The Alfvén damping rate in the isotropic case is given by

$$S_{RH,LH}(A_p = A_e = 1) = -\frac{\pi^{1/2} b^2}{2(1+\mu)\Theta} \left[\frac{1}{\mu^{1/2}} e^{-\frac{\mu}{\Theta^2} (R \pm \frac{b}{\mu})^2} + e^{-\frac{(R \mp b)^2}{\Theta}} \right] \quad (40)$$

In order to drive the Alfvén modes unstable, the growth rate (39) has to be positive requiring that

$$I_{RH,LH}(A_p, A_e, \beta_{\parallel}) \equiv \pm \left(\frac{A_p - 1}{\mu^{3/2}} e^{-\frac{w^2}{\mu \beta_{\parallel}} (1 \pm \mu x)^2} - (A_e - 1) e^{-\frac{w^2}{\beta_{\parallel}} (1 \mp x)^2} \right) + x \left[\frac{A_p}{\mu^{1/2}} e^{-\frac{w^2}{\mu \beta_{\parallel}} (1 \pm \mu x)^2} + A_e e^{-\frac{w^2}{\beta_{\parallel}} (1 \mp x)^2} \right] < 0 \quad (41)$$

This instability condition is analyzed further in the next section.

4.2. LH-polarized Alfven-proton-cyclotron and RH-polarized Alfven-Whistler-electron cyclotron branches for small parallel plasma

$$\text{beta } \beta_{\parallel} \ll 1$$

Because we are primarily interested in low plasma beta plasmas we consider for completeness the solutions of the full dispersion relation (32) for subluminal solutions $R \ll 1$ and small parallel plasma beta $\beta_{\parallel} \ll 1$:

$$0 = \Re \Lambda_{RH,LH}(x, S = 0) \simeq \frac{(1 + \mu)w^2 x^2}{1 \pm (\mu - 1)x - \mu x^2} - 1 \quad (42)$$

yielding the quadratic equation

$$[(1 + \mu)w^2 + \mu] x^2 \mp (\mu - 1)x = 1 \quad (43)$$

Because $\mu = 1836 \gg 1$ this equation is well approximated by

$$(1 + w^2)x^2 \mp x = \frac{1}{\mu} \quad (44)$$

with the solutions

$$x_{RH,LH} = \frac{1}{2(1 + w^2)} \left[\sqrt{1 + \frac{4(1 + w^2)}{\mu}} \pm 1 \right], \quad (45)$$

covering the well-known (e.g. Swanson 1989) LH-polarized Alfven-proton-cyclotron and RH-polarized Alfven-Whistler-electron cyclotron branches for parallel propagation.

For $w \ll \sqrt{(\mu/4) - 1} = 21.4$, corresponding to large wavenumbers $k \gg \omega_{p,e}/21.4c$, we obtain

$$x_{RH} \simeq \frac{1}{\mu} + \frac{1}{1 + w^2} \quad (46)$$

and

$$x_{LH} \simeq \frac{1}{\mu} \left[1 - \frac{1 + w^2}{\mu} \right] \quad (47)$$

The solution (47), subject to the constraint (24), represents the left-handed polarized proton-cyclotron waves.

For values of $w \ll 21.4$ the solution (46) reduces to

$$x_{RH} \simeq \frac{1}{1 + w^2}, \quad (48)$$

which for values of $w \ll 1$ represents the right-handed polarized electron cyclotron waves subject to the constraint (25) with the dispersion relation

$$x_{RH} \simeq 1 - w^2. \quad (49)$$

For intermediate values $1 \ll w \ll 21.4$ the solution (48) reduces to

$$x_{RH} \simeq \frac{1}{w^2}, \quad (50)$$

representing the right-handed polarized Whistler waves ($R \simeq b/w^2$).

For $w \gg \sqrt{(\mu/4) - 1} = 21.4$, corresponding to small wavenumbers $k \ll \omega_{p,e}/21.4c$, we obtain for the solution (45)

$$x_{RH,LH} \simeq \frac{1}{\sqrt{\mu(1 + w^2)}} \simeq \frac{1}{\mu^{1/2}w}, \quad (51)$$

which agrees with the Alfven wave solutions (36) in the limit $\beta_{\parallel} \ll 1$.

Eq. (42) for small plasma beta $\beta_{\parallel} \ll 1$ also provides

$$\frac{\partial \Re \Lambda_{RH,LH}(R, S=0)}{\partial R} \simeq \frac{(1+\mu)w^2x}{b} \frac{2 \pm (\mu-1)x}{[1 \pm (\mu-1)x - \mu x^2]^2} = \frac{2 \pm (\mu-1)x}{b(1+\mu)w^2x^3}, \quad (52)$$

We note that for all solutions (46) – (51), Eq. (52) is positive. According to Eqs. (17) and (26) the growth/damping rate then is

$$S_{RH,LH} = -\frac{b(1+\mu)w^2x^3}{2 \pm (\mu-1)x} \Im \Lambda_{RH,LH}(R, S=0) = -\frac{\pi^{1/2}(1+\mu)b^2w^4x^3}{\Theta[2 \pm (\mu-1)x]} \left(\frac{1}{\mu^{1/2}} \left[A_p \left[x \pm \frac{1}{\mu} \right] \mp \frac{1}{\mu} \right] e^{-\frac{\mu b^2}{\Theta^2}(x \pm \frac{1}{\mu})^2} + [A_e(x \mp 1) \pm 1] e^{-\frac{b^2(x \mp 1)^2}{\Theta^2}} \right), \quad (53)$$

which generalizes the Alfvénic growth/damping rate (39) to non-Alfvénic modes. Apart from the different notation in phase speeds, the rate (53) agrees with eq. (7.1.8) of Gary (1993). For isotropic ($A_p = A_e = 1$) plasma temperatures all modes of the RH and LH branches, including the cyclotron and Whistler modes, are damped in agreement with Brinca’s general theorem. We notice that, in order to drive the cyclotron and Whistler modes unstable, the growth rate (53) has to be positive, requiring again the earlier derived instability condition (41).

4.3. Electromagnetic light at large frequencies $R \gg b + \Theta$

For large frequencies $R \gg b + \Theta$ the dispersion relation (31) reduces to

$$\begin{aligned} 0 = \Re \Lambda_{RH,LH}(R, S=0) &\simeq R^2 - 1 - (1 + \frac{1}{\mu})w^2 \mp w^2 \frac{b}{R} - \frac{w^2 b^2}{R^2} \left[1 + \frac{1}{\mu} + \frac{\Theta^2}{2b^2} \left(A_e + \frac{A_p}{4\mu^2} \right) \right] \\ &\simeq R^2 - 1 - (1 + \frac{1}{\mu})w^2 \end{aligned} \quad (54)$$

and

$$0 = \Im \Lambda_{RH,LH}(R, S=0) \simeq \pi^{1/2}w^2 \frac{R}{\Theta} \left(\frac{A_p}{\mu^{1/2}} e^{-\mu R^2/\Theta^2} + A_e e^{-R^2/\Theta^2} \right), \quad (55)$$

so that for both polarisations to lowest order in $(b/R)^2$ we obtain the dispersion relation of electromagnetic light

$$R^2 \simeq 1 + \left(1 + \frac{1}{\mu}\right)w^2 \quad (56)$$

with the same damping rate

$$S = -\frac{\pi^{1/2}w^2}{2\Theta} \left(\frac{A_p}{\mu^{1/2}} e^{-\mu R^2/\Theta^2} + A_e e^{-R^2/\Theta^2} \right) \quad (57)$$

5. Instability conditions

In order to drive RH-handed polarized Alfven-proton cyclotron and the LH-handed polarized Alfven-Whistler-electron cyclotron branches unstable, the condition (41) has to be fulfilled which, after multiplication with $\mu^{3/4}$, reads

$$\begin{aligned} & \pm \left(\frac{A_p - 1}{\mu^{3/4}} e^{-X_p^2} - (A_e - 1) \mu^{3/4} e^{-X_e^2} \right) + x \mu^{1/2} \left[\frac{A_p}{\mu^{1/4}} e^{-X_p^2} + A_e \mu^{1/4} e^{-X_e^2} \right] = \\ & \pm \left(e^{-\frac{3}{4} \ln \mu + \ln(A_p - 1) - X_p^2} - e^{\frac{3}{4} \ln \mu + \ln(A_e - 1) - X_e^2} \right) \\ & + x \mu^{1/2} \left(e^{-\frac{1}{4} \ln \mu + \ln A_p - X_p^2} + e^{\frac{1}{4} \ln \mu + \ln A_e - X_e^2} \right) < 0 \end{aligned} \quad (58)$$

with

$$X_p = \frac{w}{(\mu \beta_{\parallel})^{1/2}} (1 \pm \mu x), \quad X_e = \frac{w}{\beta_{\parallel}^{1/2}} (1 \mp x). \quad (59)$$

Both brackets in Eq. (58) can be further reduced using the identities

$$\begin{aligned} e^{a_1+a_2} + e^{a_1-a_2} &= 2e^{a_1} \cosh(a_2), \quad e^{a_1+a_2} - e^{a_1-a_2} = 2e^{a_1} \sinh(a_2), \\ e^{c_1+c_2} + e^{c_1-c_2} &= 2e^{c_1} \cosh(c_2), \quad e^{c_1+c_2} - e^{c_1-c_2} = 2e^{c_1} \sinh(c_2) \end{aligned} \quad (60)$$

From the first bracket we identify

$$a_1 = \ln [(A_p - 1)(A_e - 1)]^{1/2} - \frac{X_p^2 + X_e^2}{2}, \quad a_2 = -\frac{3}{4} \ln \mu - \ln \left[\frac{A_e - 1}{A_p - 1} \right]^{1/2} - W_{\pm} \quad (61)$$

where

$$W_{\pm}(x) = \frac{X_p^2 - X_e^2}{2} = \frac{w^2}{2\beta_{\parallel}} \left[\pm 4x + (\mu - 1)(x^2 - \frac{1}{\mu}) \right], \quad (62)$$

From the second bracket we determine

$$c_1 = \ln [A_p A_e]^{1/2} - \frac{X_p^2 + X_e^2}{2}, \quad c_2 = -\frac{1}{4} \ln \mu - \ln \left[\frac{A_e}{A_p} \right]^{1/2} - W_{\pm} \quad (63)$$

With $\sinh(-a_2) = -\sinh(a_2)$ and $\cosh(-c_2) = \cosh(c_2)$ condition (58) becomes

$$\begin{aligned} & \pm (A_p - 1)^{1/2} (A_e - 1)^{1/2} \sinh \left[\frac{3}{4} \ln \mu + \ln \left[\frac{A_e - 1}{A_p - 1} \right]^{1/2} + W_{\pm} \right] \\ & - A_p^{1/2} A_e^{1/2} x \mu^{1/2} \cosh \left[\frac{1}{4} \ln \mu + \ln \left[\frac{A_e}{A_p} \right]^{1/2} + W_{\pm} \right] \\ & = \pm (\mu^{3/4} (A_e - 1) - \mu^{-3/4} (A_p - 1)) \cosh W_{\pm} \pm (\mu^{3/4} (A_e - 1) + \mu^{-3/4} (A_p - 1)) \sinh W_{\pm} \\ & - x \mu^{1/2} (\mu^{1/4} A_e + \mu^{-1/4} A_p) \cosh W_{\pm} - x \mu^{1/2} (\mu^{1/4} A_e - \mu^{-1/4} A_p) \sinh W_{\pm} > 0, \end{aligned} \quad (64)$$

which yields the general instability condition for the two branches in the form

$$\begin{aligned} & \pm \frac{(A_e - A_p)[(\mu^{3/2} + 1) + (\mu^{3/2} - 1) \tanh(W_{\pm})] + (A_e + A_p - 2)[(\mu^{3/2} - 1) + (\mu^{3/2} + 1) \tanh(W_{\pm})]}{x \mu [(A_e + A_p)[\mu^{1/2} + 1 + (\mu^{1/2} - 1) \tanh(W_{\pm})] + (A_e - A_p)[\mu^{1/2} - 1 + (\mu^{1/2} + 1) \tanh(W_{\pm})]} \\ & > 1 \end{aligned} \quad (65)$$

Inserting the general dispersion relation for small plasma betas (45) for x and $W_{\pm}(x)$ then provides the general instability conditions for the RH-handed polarized Alfvén-proton

cyclotron and the LH-handed polarized Alfvén-Whistler-electron cyclotron branches, which to the best of our knowledge has not been derived before.

For equal electron and proton temperature anisotropies ($A_p = A_e = A_0$) the condition (65) reduces to

$$\begin{aligned} & \pm \left(1 - \frac{1}{A_0}\right) [(\mu^{3/2} - 1) + (\mu^{3/2} + 1) \tanh(W_{\pm})] \\ & > \mu x [\mu^{1/2} + 1 + (\mu^{1/2} - 1) \tanh(W_{\pm})]. \end{aligned} \quad (66)$$

Note that for pair plasmas ($\mu = 1$) the condition (66) reduces to Eq. (S-66) in paper S.

Instead of working with the general dispersion relations (45) used in the general instability condition (65), we will discuss different ranges of the plasma frequency phase speed w , where the dispersion relations (45) reduce to simpler limits, as demonstrated in Sect. 4.2. For $w \gg 21.4$ we can use the Alfvénic relation (51), whereas for small values of $w < 21.4$ we can use the cyclotron and Whistler relations (47), (48), (49) and (50), respectively.

Before investigating the individual modes, we inspect the instability condition (66). The instability condition holds for plasma betas $\beta_{\parallel} \ll w^2/\mu$, implying $w^2/2\beta_{\parallel} \gg \mu/2 = 918$. Consequently, the argument of the tanh-function

$$W_{\pm} = \frac{w^2}{2\beta_{\parallel}} f_{\pm}(x), \quad f_{\pm}(x) = \pm 4x + (\mu - 1)(x^2 - \frac{1}{\mu}) \quad (67)$$

is much larger than unity for $f_{\pm}(x) \gg 2/\mu$.

The function $f_+(x)$ increases monotonically from its smallest negative value $f_+(0) = -(1 - \mu^{-1}) = -0.9995$ and becomes zero at

$$x_{0+} = \frac{\mu + 1}{(\mu - 1)\mu^{1/2}} \left[1 - \frac{2\mu^{1/2}}{\mu + 1}\right] = 0.95\mu^{-1/2} \simeq \mu^{-1/2} \quad (68)$$

Likewise, the function $f_-(x)$ attains its negative minimum value -1.0016 at $x_E = 2/(\mu - 1) = 0.0011$ and becomes zero at

$$x_{0-} = \frac{\mu + 1}{(\mu - 1)\mu^{1/2}} \left[1 + \frac{2\mu^{1/2}}{\mu + 1} \right] = 1.05\mu^{-1/2} \simeq \mu^{-1/2} \quad (69)$$

Hence only for values of $x = R/b$ inside a small interval around $\mu^{-1/2}$, which for right-handed polarized fluctuations lies in the Whistler phase speed range, the functions $f_{\pm}(x)$ are smaller than $2/\mu$. Both functions are well approximated by

$$f_{\pm}(x) \simeq f_0(x) = \mu x^2 - 1, \quad (70)$$

which is negative in the range of Alfven and proton cyclotron waves and positive in the range of electron cyclotron waves, respectively.

5.1. Left-handed polarized Alfven waves and proton cyclotron waves

In the range of Alfven and proton cyclotron waves $x \leq \mu^{-1}$ the function $f_-(x) \simeq -1$ is negative and practically constant, so that the instability condition (66) becomes

$$\begin{aligned} & \left(1 - \frac{1}{A_0} \right) \left[(\mu^{3/2} + 1) \tanh \left[\frac{w^2}{2\beta_{\parallel}} \right] - (\mu^{3/2} - 1) \right] \\ & > x\mu \left[(\mu^{1/2} + 1) - (\mu^{1/2} - 1) \tanh \left[\frac{w^2}{2\beta_{\parallel}} \right] \right] \end{aligned} \quad (71)$$

The tanh-function is well approximated by

$$\tanh(t) \simeq \frac{t}{1 + t}, \quad (72)$$

so that with $2\beta_{\parallel}/w^2 \ll 2/\mu = 1/918 \ll 1$

$$\tanh \left[\frac{w^2}{2\beta_{\parallel}} \right] \simeq \frac{1}{1 + \frac{2\beta_{\parallel}}{w^2}} \simeq 1 - \frac{2\beta_{\parallel}}{w^2}, \quad (73)$$

we find for condition (71)

$$\left(1 - \frac{1}{A_0} \right) \left[1 - (1 + \mu^{3/2}) \frac{\beta_{\parallel}}{w^2} \right] > \mu x \left[1 + (\mu^{1/2} - 1) \frac{\beta_{\parallel}}{w^2} \right]. \quad (74)$$

Because $\mu^{3/2} \gg 1$ and $\mu^{1/2} \gg 1$ condition (74) is well approximated by

$$\left(1 - \frac{1}{A_0} \right) [1 - \eta] > x[\mu + \eta], \quad (75)$$

where

$$\eta = \frac{\beta_{\parallel}(\mu^{3/2} + 1)}{w^2} = \beta_{\parallel}/\beta_c \quad (76)$$

denotes a normalized parallel plasma beta value in terms of the critical plasma beta

$$\beta_c = \frac{w^2}{(1 + \mu^{3/2})} \quad (77)$$

The restriction $\beta_{\parallel} \ll w^2/\mu$ requires $\eta \ll \mu^{1/2} = 43$. Obviously, we have to consider the two cases $\eta < 1$ and $1 < \eta < 43$, corresponding to $\beta_{\parallel} < \beta_c$ and $\beta_c < \beta_{\parallel} < 43\beta_c$, respectively.

For $\beta_{\parallel} < \beta_c$ the instability condition (75) becomes

$$1 - \frac{1}{A_0} > x \frac{\mu + \eta}{1 - \eta}, \quad (78)$$

or

$$A_0(\beta_{\parallel} < \beta_c) > \frac{1 - \eta}{(1 - \mu x) - (1 + x)\eta} \quad (79)$$

Likewise, for $\beta_c < \beta_{\parallel} < 43\beta_c$ we obtain

$$\left(\frac{1}{A_0} - 1\right) > x \frac{\mu + \eta}{\eta - 1}, \quad (80)$$

corresponding to

$$A_0(\beta_{\parallel} > \beta_c) < \frac{\eta - 1}{(1 + x)\eta - (1 - \mu x)} \quad (81)$$

For plasma beta values equal to the critical value ($\eta = 1$), the instability condition (75) cannot be fulfilled. The two instability conditions (79) and (81) for the LH polarized Alfvén and proton cyclotron waves are illustrated in Fig. 2 together with the instability conditions for the RH polarized Alfvén and low phase speed Whistler waves, which we derive in the next subsection.

We remark at this point that the threshold condition for the LH Alfvén-proton cyclotron branch shown in Fig. 2 does not agree with the AIC threshold condition plotted in Fig. 1 of Bale et al. (2009).

5.2. Right-handed polarized Alfvén waves and low phase speed Whistler waves

In the range of the RH polarized Alfvén and Whistler waves below $x \ll \mu^{-1/2}$ the function $f_+(x) \simeq -1$ is negative and practically constant, so that the instability condition (66) together with $W_+ = -w^2/2\beta_{\parallel}$ becomes

$$\begin{aligned} & \left(\frac{1}{A_0} - 1\right) \left[(\mu^{3/2} + 1) \tanh \left[\frac{w^2}{2\beta_{\parallel}} \right] - (\mu^{3/2} - 1) \right] \\ & > x\mu \left[(\mu^{1/2} + 1) - (\mu^{1/2} - 1) \tanh \left[\frac{w^2}{2\beta_{\parallel}} \right] \right] \end{aligned} \quad (82)$$

With the approximation (72) we derive for condition (82)

$$\left(\frac{1}{A_0} - 1\right)(1 - \eta) > x(\mu + \eta) \quad (83)$$

For $\beta_{\parallel} < \beta_c$ the instability condition (83) becomes

$$A_0(\beta_{\parallel} < \beta_c) < \frac{1 - \eta}{(1 + \mu x) - (1 - x)\eta} \quad (84)$$

Likewise, for $\beta_c < \beta_{\parallel} < 43\beta_c$ we obtain

$$A_0(\beta_{\parallel} > \beta_c) > \frac{\eta - 1}{(1 - x)\eta - (1 + \mu x)} \quad (85)$$

For plasma beta values equal to the critical value ($\eta = 1$), the instability condition (75) cannot be fulfilled. The two instability conditions (84) and (85) for the RH polarized Alfvén and low phase speed Whistler waves are illustrated in Fig. 2.

5.3. Right-handed polarized high phase speed Whistler waves and electron cyclotron waves

In the range of high speed Whistler and electron cyclotron waves $x \gg \mu^{-1/2}$ the function $f_+(x) \simeq \mu x^2$ is positive. The instability condition (66) together with $W_+ = w^2 f_+(x)/(2\beta_{\parallel}) = \mu w^2 x^2/(2\beta_{\parallel})$ then becomes

$$\begin{aligned} \left(1 - \frac{1}{A_0}\right) \left[(\mu^{3/2} - 1) + (\mu^{3/2} + 1) \tanh \left[\frac{\mu w^2 x^2}{2\beta_{\parallel}} \right] \right] > \\ x\mu \left[(\mu^{1/2} + 1) + (\mu^{1/2} - 1) \tanh \left[\frac{\mu w^2 x^2}{2\beta_{\parallel}} \right] \right] \end{aligned} \quad (86)$$

With the approximation (72) we derive for condition (86)

$$\left(1 - \frac{1}{A_0}\right) \left[1 - \frac{\beta_{\parallel}}{\mu w^2 x^2} \right] > x \left[1 + \frac{\beta_{\parallel}}{\mu w^2 x^2} \right]. \quad (87)$$

Because $x \gg \mu^{-1/2}$ we find that

$$\frac{\beta_{\parallel}}{\mu w^2 x^2} \ll \frac{\beta_{\parallel}}{w^2} \ll \mu^{-1}, \quad (88)$$

so that

$$1 - \frac{1}{A_0} > x \frac{1 + \frac{\beta_{\parallel}}{\mu w^2 x^2}}{1 - \frac{\beta_{\parallel}}{\mu w^2 x^2}} \simeq x \left[1 + \frac{2\beta_{\parallel}}{\mu w^2 x^2} \right], \quad (89)$$

leading to

$$A_0 > \frac{1}{1 - x \left[1 + \frac{2\beta_{\parallel}}{\mu w^2 x^2} \right]} \quad (90)$$

We now discuss the instability conditions for individual wave modes.

6. RH and LH polarized Alfven waves

For $w \gg \mu^{1/2} > 21.4$ we insert the Alfven wave dispersion relation (51) in the conditions (79), (81), (84) and (85). For $\beta_{\parallel} < \beta_c$, corresponding to $\eta < 1$, we obtain for the RH polarized Alfven waves

$$A_{0,RH}(\beta_{\parallel} < \beta_c) < \frac{w(1 - \eta)}{(w + \mu^{1/2}) - (w - \mu^{-1/2})\eta} = \frac{w(\beta_c - \beta_{\parallel})}{(w + \mu^{1/2})\beta_c - (w - \mu^{-1/2})\beta_{\parallel}} \quad (91)$$

and the LH polarized Alfven waves

$$A_{0,LH}(\beta_{\parallel} < \beta_c) > \frac{w(1 - \eta)}{(w - \mu^{1/2}) - (w + \mu^{-1/2})\eta} = \frac{w(\beta_c - \beta_{\parallel})}{(w - \mu^{1/2})\beta_c - (w + \mu^{-1/2})\beta_{\parallel}} \quad (92)$$

Likewise, for $\beta_{\parallel} > \beta_c$, corresponding to $\eta > 1$, we obtain for the RH polarized Alfven waves

$$A_{0,RH}(\beta_{\parallel} > \beta_c) > \frac{w(\eta - 1)}{(w - \mu^{1/2})\eta - (w + \mu^{-1/2})} = \frac{w(\beta_{\parallel} - \beta_c)}{(w - \mu^{1/2})\beta_{\parallel} - (w + \mu^{-1/2})\beta_c} \quad (93)$$

and the LH polarized Alfven waves

$$A_{0,LH}(\beta_{\parallel} > \beta_c) < \frac{w(\eta - 1)}{(w + \mu^{-1/2})\eta - (w - \mu^{1/2})} = \frac{w(\beta_{\parallel} - \beta_c)}{(w + \mu^{-1/2})\beta_{\parallel} - (w - \mu^{1/2})\beta_c} \quad (94)$$

These four Alfvenic instability conditions are shown in Fig. 3. Note, as indicated in the figure, that the limiting anisotropy values and parallel plasma betas for $\beta_{\parallel} \rightarrow 0$ and $\beta_{\parallel} \rightarrow \infty$ and $A_0 \rightarrow \infty$, respectively, are solely determined by the mass ratio μ and the wavenumber-dependent plasma frequency phase speed $w = \omega_{p,e}/kc$.

The properties of the weakly amplified polarized LH and RH polarized Alfven mode are summarized in Table 1 and 2, respectively.

7. Proton cyclotron waves

For $w \ll 21.4$ we insert the dispersion relation (47) for proton cyclotron waves in the instability conditions (79) and (81) yielding

$$A_0(\beta_{\parallel} < \beta_c) > \frac{\mu(1 - \eta)}{1 + w^2 - (\mu + 1)\eta} = \frac{\mu(\beta_c - \beta_{\parallel})}{(1 + w^2)\beta_c - (\mu + 1)\beta_{\parallel}} \quad (95)$$

and

$$A_0(\beta_{\parallel} > \beta_c) < \frac{\mu(\eta - 1)}{(\mu + 1)\eta - (1 + w^2)} = \frac{\mu(\beta_{\parallel} - \beta_c)}{(\mu + 1)\beta_{\parallel} - (1 + w^2)\beta_c} \quad (96)$$

These instability conditions cannot be fulfilled for anisotropies in the interval $A_0 \in [\mu/(1 + \mu), \mu/(1 + w^2)]$, including the isotropic case $A_0 = 1$. The proton cyclotron instability conditions are illustrated in Fig. 4 as a function of the parallel plasma beta $\beta_{\parallel}$.

Real phase speed range	$R \ll b/\mu = w\Theta/(\mu\beta_{\parallel}^{1/2})$
Parallel plasma beta range	$\Theta^2 \ll \beta_{\parallel} \ll w^2/\mu$
Dispersion relation	$x = R/b = \frac{1}{\mu^{1/2}w}$
Existence condition	$w \gg \mu^{1/2} = 43 > \sqrt{(\mu/4) - 1} = 21.4$
Instability conditions:	$A_0 \notin \left[\frac{w}{w+\mu^{-1/2}}, \frac{w}{w-\mu^{1/2}} \right] = \left[\frac{43w}{1+43w}, \frac{w}{w-43} \right]$
for $\beta_{\parallel} < \beta_c = \frac{w^2}{(1+\mu^{3/2})}$	$A_0 > \frac{w(\beta_c - \beta_{\parallel})}{(w-\mu^{1/2})\beta_c - (w+\mu^{-1/2})\beta_{\parallel}}$
for $\beta_{\parallel} > \beta_c$	$A_0 < \frac{w(\beta_{\parallel} - \beta_c)}{(w+\mu^{-1/2})\beta_{\parallel} - (w-\mu^{1/2})\beta_c}$

Table 1: Properties of weakly amplified LH polarized Alfven wave mode

Real phase speed range	$R \ll b/\mu = w\Theta/(\mu\beta_{\parallel}^{1/2})$
Parallel plasma beta range	$\Theta^2 \ll \beta_{\parallel} \ll w^2/\mu$
Dispersion relation	$x = R/b = \frac{1}{\mu^{1/2}w}$
Existence condition	$w \gg \mu^{1/2} = 43 > \sqrt{(\mu/4) - 1} = 21.4$
Instability conditions:	$A_0 \notin \left[\frac{w}{w+\mu^{1/2}}, \frac{w}{w-\mu^{-1/2}} \right] = \left[\frac{w}{w+43}, \frac{43w}{43w-1} \right]$
for $\beta_{\parallel} < \beta_c = \frac{w^2}{(1+\mu^{3/2})}$	$A_0 < \frac{w(\beta_c - \beta_{\parallel})}{(w+\mu^{1/2})\beta_c - (w-\mu^{-1/2})\beta_{\parallel}}$
for $\beta_{\parallel} > \beta_c$	$A_0 > \frac{w(\beta_{\parallel} - \beta_c)}{(w-\mu^{1/2})\beta_{\parallel} - (w+\mu^{-1/2})\beta_c}$

Table 2: Properties of weakly amplified RH polarized Alfven wave mode

The properties of the weakly amplified proton cyclotron mode are summarized in Table 3.

8. Right-handed polarized low phase speed Whistler waves

The Whistler-electron cyclotron dispersion relation (48) yields values of $x \ll \mu^{-1/2}$ provided

$$\frac{1}{1+w^2} \ll \mu^{-1/2}, \quad (97)$$

which is equivalent to

$$(\mu^{1/2} - 1)^{1/2} = 6.5 \ll w \ll 21.4 \quad (98)$$

The instability conditions (84) and (85) then reduce to

$$A_0(\beta_{\parallel} < \beta_c) < \frac{(1+w^2)(1-\eta)}{(1+\mu+w^2)-w^2\eta} = \frac{(1+w^2)(\beta_c - \beta_{\parallel})}{(1+\mu+w^2)\beta_c - w^2\beta_{\parallel}} \quad (99)$$

and

$$A_0(\beta_{\parallel} > \beta_c) > \frac{(1+w^2)(\eta-1)}{w^2\eta - (1+\mu+w^2)} = \frac{(1+w^2)(\beta_{\parallel} - \beta_c)}{w^2\beta_{\parallel} - (1+\mu+w^2)\beta_c} \quad (100)$$

These instability condition cannot be fulfilled for anisotropies in the interval

$$\frac{1+w^2}{1+w^2+\mu} \leq A_0 \leq \frac{1+w^2}{w^2}, \quad (101)$$

including the isotropic case $A_0 = 1$. The RH low phase speed Whistler wave instability conditions are illustrated in Fig. 5 as a function of the parallel plasma beta $\beta_{\parallel}$.

The properties of the weakly amplified RH polarized low phase speed Whistler mode are summarized in Table 4.

9. Right-handed polarized electron cyclotron waves and high phase speed Whistler waves

The Whistler-electron cyclotron dispersion relation (48) yields values of $x \gg \mu^{-1/2}$ provided

$$\frac{1}{1+w^2} \gg \mu^{-1/2}, \quad (102)$$

which is equivalent to

$$w \ll (\mu^{1/2} - 1)^{1/2} = 6.5 \quad (103)$$

Inserting the Whistler-electron cyclotron dispersion relation (48) yields for the instability condition (90)

$$A_0 > \frac{\mu w^2(1+w^2)}{\mu w^4 - 2(1+w^2)^2 \beta_{\parallel}}, \quad (104)$$

which is illustrated in Fig. 6.

The properties of the weakly amplified RH polarized high phase speed Whistler and electron cyclotron modes are summarized in Table 5.

Real phase speed range	$R \ll b$
Parallel plasma beta range	$\Theta^2 \ll \beta_{\parallel} \ll \frac{w^2}{\mu}$
Dispersion relation	$x = R/b = \frac{1}{\mu} [1 - \frac{1+w^2}{\mu}]$
Existence condition	$w \ll \sqrt{(\mu/4) - 1} = 21.4$
Instability conditions:	$A_0 \notin \left[\frac{\mu}{1+\mu}, \frac{\mu}{1+w^2} \right]$
for $\beta_{\parallel} < \beta_c = \frac{w^2}{(1+\mu^{3/2})}$	$A_0 > \frac{\mu(\beta_c - \beta_{\parallel})}{(1+w^2)\beta_c - (\mu+1)\beta_{\parallel}}$
for $\beta_{\parallel} > \beta_c$	$A_0 < \frac{\mu(\beta_{\parallel} - \beta_c)}{(\mu+1)\beta_{\parallel} - (1+w^2)\beta_c}$

Table 3: Properties of weakly amplified LH polarized proton cyclotron mode

Real phase speed range	$\frac{b}{\mu} \ll R \ll b\mu^{-1/2}$
Parallel plasma beta range	$\Theta^2 \ll \beta_{\parallel} \ll w^2/\mu$
Dispersion relation	$x = R/b = \frac{1}{1+w^2}$
Existence condition	$6.5 \ll w \ll 21.4$
Instability conditions:	$A_0 \notin \left[\frac{1+w^2}{1+\mu+w^2}, \frac{1+w^2}{w^2} \right]$
for $\beta_{\parallel} < \beta_c = \frac{w^2}{(1+\mu^{3/2})}$	$A_0 < \frac{(1+w^2)(\beta_c - \beta_{\parallel})}{(1+\mu+w^2)\beta_c - w^2\beta_{\parallel}}$
for $\beta_{\parallel} > \beta_c$	$A_0 > \frac{(1+w^2)(\beta_{\parallel} - \beta_c)}{w^2\beta_{\parallel} - (1+\mu+w^2)\beta_c}$

Table 4: Properties of weakly amplified RH polarized low phase speed Whistler mode

Real phase speed range	$b\mu^{-1/2} \ll R \ll b$
Parallel plasma beta range	$\Theta^2 \ll \beta_{\parallel} \ll w^2/\mu$
Dispersion relation	$x = R/b = \frac{1}{1+w^2}$
Existence condition	$w \ll 6.5$
Instability condition:	
for $\beta_{\parallel} < w^2/\mu$	$A_0 > \frac{\mu w^2(1+w^2)}{\mu w^4 - 2(1+w^2)^2\beta_{\parallel}}$

Table 5: Properties of weakly amplified RH polarized high phase speed Whistler and electron cyclotron wave mode

10. Application to solar wind fluctuations

The solar wind magnetic fluctuations measured by Bale et al. (2009) near 1 AU have wavenumbers

$$k \simeq \alpha / \rho_p \quad (105)$$

with $\alpha = 0.56 \pm 0.32$ and the thermal proton gyroradius $\rho_p = 4.23 \cdot 10^6 T_5^{1/2} B_4^{-1}$ cm, where we adopt an interplanetary magnetic field value $B = 10^{-4} B_4$ Gauss and a temperature $T = 10^5 T_5$ K. With the solar wind particle density $n_e = 10^2 n_2$ cm⁻³ we find that the plasma frequency phase speed (8)

$$w = \frac{79.5}{\alpha} \frac{(T_5 n_2)^{1/2}}{B_4} = \frac{142}{1 \pm 0.57} \frac{(T_5 n_2)^{1/2}}{B_4} \quad (106)$$

covers the interval

$$90 \frac{(T_5 n_2)^{1/2}}{B_4} \leq w \leq 330 \frac{(T_5 n_2)^{1/2}}{B_4} \quad (107)$$

The electron gyrofrequency phase speed (9)

$$b = 3.13 \cdot 10^{-3} w \frac{B_4}{n_2^{1/2}} \quad (108)$$

is indeed much smaller than the electron plasma frequency phase speed w justifying the high density plasma approximation $w \gg b$.

According to the existence conditions of the different instabilities for low plasma beta values summarized in Tables 1-5, only the left-handed and right-handed polarized Alfvén wave instabilities can operate in the w -interval (107). Fig. 3 therefore applies to this application. Fig. 3 was calculated for a value of $w = 280$ lying within the range

(107) of the observed solar wind turbulence. $w = 280$ yields for the critical plasma beta $\beta_c = w^2/(1 + \mu^{3/2}) = 0.997$.

Besides the mass ratio μ the Alfvénic instability threshold diagram shown in Fig. 3 is controlled by the single observed plasma parameter w . These two quantities define the characteristic plasma beta $\beta_c = w^2/\mu^{3/2}$ and the parameter

$$a = \frac{\mu^{1/2} - \mu^{-1/2}}{w - \mu^{-1/2}} \simeq \frac{\mu^{1/2}}{w} \quad (109)$$

As illustrated in Fig. 7 all asymptotic and characteristic values of the threshold anisotropies and the characteristic plasma beta values in the case of LH and RH polarized Alfvén waves are solely determined by a and β_c .

11. Summary and conclusions

We rigorously studied the dispersion relations of weakly amplified fluctuations with wave vectors $\vec{k} \times \vec{B}_0 = 0$ in an anisotropic bi-Maxwellian magnetized proton-electron plasma by the appropriate Taylor expansion of the plasma dispersion function. Apparently for the first time, for equal parallel electron and proton temperatures, a general analytical instability condition (65) is derived that holds for different values of the electron (A_e) and proton (A_p) temperature anisotropies. We determine the conditions for which the weakly amplified LH-handed polarized Alfvén-proton-cyclotron and RH-handed polarized Alfvén-Whistler-electron-cyclotron branches can be excited. For different regimes of the electron plasma frequency phase speed $w = \omega_{p,e}/(kc)$ these branches reduce to the RH and LH polarized Alfvén waves, RH polarized high- and low-phase speed Whistler, RH polarized proton and LH polarized electron cyclotron modes. The properties of the individual modes are summarized in Tables 1-5. Analytic instability threshold conditions are derived

in terms of the combined temperature anisotropy $A = T_{\perp}/T_{\parallel}$, the parallel plasma beta $\beta_{\parallel} = 8\pi n_e k_B T_{\parallel}/B^2$ and the electron plasma frequency phase speed $w = \omega_{p,e}/(kc)$ for each mode.

For large values of $w \gg 21.4$, corresponding to small wavenumbers $k \ll \omega_{p,e}/21.4c$, RH and LH polarized Alfvén waves are excited in different regions of the temperature anisotropy (A)-versus parallel plasma beta ($\beta_{\parallel}$)-parameter plane. For appropriate temperature anisotropies, RH polarized low and high phase speed Whistler waves are excited for intermediate values of $6.5 \ll w \ll 21.4$ and $1 \ll w \ll 6.5$, respectively. At values $w \ll 21.4$ LH polarized proton cyclotron waves can be excited, whereas RH polarized electron cyclotron waves can be excited at small values of $w \leq 1$. In agreement with the general theorem of Brinca (1990) on the electromagnetic stability of isotropic plasma populations none of these modes can be excited for isotropic plasma distributions ($A_e = A_p = 1$).

We apply the results of our instability study to the observed solar wind magnetic turbulence (Bale et al. 2009), corresponding to values of $90 \leq w \leq 330$. According to the existence conditions of the different instabilities, only the left-handed and right-handed polarized Alfvén wave instabilities can operate here. Besides the electron-proton mass ratio $\mu = 1836$, the Alfvénic instability threshold conditions are controlled by the single observed plasma parameter w . All asymptotic and characteristic values of the threshold anisotropies and the characteristic plasma beta values are solely determined by the values of $a = 43/w$ and $\beta_c = w^2/\mu^{3/2}$. Comparing the Alfvénic instability diagram shown in Fig. 7 with the observations shown in Fig. 1 of Bale et al. (2009), one notices that the main characteristic properties of the observed solar wind fluctuations are well reproduced by the instability conditions of parallel propagating Alfvén waves. Especially, the observed confinement limits at small parallel plasma beta values are explained. It remains to be investigated in future work how weakly propagating and/or obliquely propagating weakly amplified modes add to

this diagram.

We thank the referee for his constructive comments that helped to improve the manuscript. This work was supported by the Deutsche Forschungsgemeinschaft through grants Schl 201/19-1 and Schl 201/21-1.




Fig. 2.— Anisotropy diagram for LH polarized Alfvén and proton cyclotron waves and for RH polarized Alfvén and low phase speed Whistler waves for the case of equal electron and proton temperature anisotropies $A_e = A_p = A_0$. $\eta = \beta_{\parallel}/(w^2/\mu)$ denotes the normalized plasma beta. Unstable regions are marked by "u",




Fig. 3.— Anisotropy diagram for LH and RH polarized Alfvén waves for the case of equal electron and proton temperature anisotropies $A_e = A_p = A_0$. Stable regions are marked by "S", unstable regions are marked by "u". A value of $w = 280$ is adopted so that the results hold for parallel plasma beta values $\beta_{\parallel} < 5.4$.




Fig. 4.— Anisotropy diagram for LH polarized proton cyclotron waves for the case of equal electron and proton temperature anisotropies $A_e = A_p = A_0$. Unstable regions are marked by "u", A value of $w = 10$ is adopted so that the results hold for parallel plasma beta values $\beta_{\parallel} < 0.054$.




Fig. 5.— Anisotropy diagram for RH polarized low phase speed Whistler waves for the case of equal electron and proton temperature anisotropies $A_e = A_p = A_0$. Unstable regions are marked by "u". A value of $w = 10$ is adopted so that the results hold for parallel plasma beta values $\beta_{\parallel} < 0.054$.




Fig. 6.— Anisotropy diagram for RH polarized high phase speed Whistler and electron cyclotron waves for the case of equal electron and proton temperature anisotropies $A_e = A_p = A_0$. Unstable regions are marked by "u", Values of $w = 3$ and $w = 0.01$ are adopted for high phase speed Whistlers and electron cyclotron waves, respectively, so that the results hold for parallel plasma beta values below $5 \cdot 10^{-3}$ and $5 \cdot 10^{-6}$, respectively.




Fig. 7.— Anisotropy diagram for LH and RH polarized Alfvén waves in terms of the plasma parameters β_c and $a = \mu^{1/2}/w$. Unstable regions are marked by "u",

REFERENCES

- Bale, S. D., Kasper, J. C., Howes, G. G., et al. 2009, *Physical Review Letters*, 103, 211101
- Brinca, A. L. 1990, *J. Geophys. Res.*, 95, 221
- Cuperman, S. 1992, *Rev. Geophys. Space Phys.*, 19, 307
- Fried, B. D. & Conte, S. D. 1961, *The Plasma Dispersion Function* ()
- Gary, S. P. 1993, *Theory of Space Plasma Microinstabilities* ()
- Gary, S. P. & Lee, M. A. 1994, *J. Geophys. Res.*, 99, 11297
- Gary, S. P., McKean, M. E., Winske, D., et al. 1994, *J. Geophys. Res.*, 99, 5903
- Gary, S. P., Skoug, R. M., Steinberg, J. T., & Smith, C. W. 2001, *Geophys. Res. Lett.*, 28, 2759
- Hellinger, P., Trávníček, P., Kasper, J. C., & Lazarus, A. J. 2006, *Geophys. Res. Lett.*, 33, 9101
- Kasper, J. C., Lazarus, A. J., & Gary, S. P. 2002, *Geophys. Res. Lett.*, 29, 170000
- Marsch, E. 2006, *Living Reviews in Solar Physics*, 3, 1
- Pokhotelov, O. A., Sagdeev, R. Z., Balikhin, M. A., & Treumann, R. A. 2004, *Journal of Geophysical Research (Space Physics)*, 109, 9213
- Samsonov, A. A., Pudovkin, M. I., Gary, S. P., & Hubert, D. 2001, *J. Geophys. Res.*, 106, 21689
- Schekochihin, A. A., Cowley, S. C., Kulsrud, R. M., Hammett, G. W., & Sharma, P. 2005, *ApJ*, 629, 139

Schlickeiser, R. 2010, The Open Plasma Physics Journal, 3, 1

Sharma, P., Quataert, E., Hammett, G. W., & Stone, J. M. 2007, ApJ, 667, 714

Swanson, D. G. 1989, Plasma waves. ()